

KINETICS OF SUB-CRITICAL BUBBLES AND THE ELECTROWEAK TRANSITIONGraciela Gelmini^a and Marcelo Gleiser^b*^a Department of Physics,
University of California, Los Angeles, CA 90024**^b Department of Physics and Astronomy,
Dartmouth College, Hanover, NH 03755***ABSTRACT**

We investigate the role of large amplitude sub-critical thermal fluctuations in the dynamics of first order phase transitions. In particular, we obtain a kinetic equation for the number density of sub-critical fluctuations of the broken-symmetric phase within the symmetric phase, modeled as spherical bubbles, and solve it analytically for temperatures above the critical temperature. We study the approach to equilibrium and obtain the equilibrium distribution of sub-critical bubbles of the unstable phase by examining three possible mechanisms responsible for their removal; their shrinking, their coupling to thermal noise, and by thermal fluctuations of the true vacuum inside them. We show that for sufficiently strong transitions, either the shrinking or the coupling to thermal noise dominate the dynamics. As the strength of the transition weakens we show that sub-critical fluctuations become progressively more important, as a larger fraction of the total volume is occupied by the broken-symmetric phase, until the point where our analytical approach breaks down. Our investigation suggests that pre-transitional phenomena may considerably change the dynamics of sufficiently weak first-order transitions. We apply our results to the standard electroweak transition.

e-mail: gelmini@physics.ucla.edu; gleiser@peterpan.dartmouth.edu

1. INTRODUCTION

The study of the kinetics of first order phase transitions is by no means a new topic. Since the pioneer work of Becker and Döring, the condensation of supersaturated vapor has been understood to occur by the thermal nucleation of droplets of the liquid phase which, when larger than a critical size, will grow and coalesce completing the transition.¹ Despite all the effort dedicated to the study of the kinetics of phase transitions, their intrinsic non-linear nature makes progress slow and restricted to simple systems. Even for simple systems in the same universality class of the Ising model many questions remain unanswered, from a microscopic understanding of coarse-graining to the description of the late stages of the transition.²

Our main interest is in the study of cosmological first order transitions, where the cooling is enforced adiabatically by the expanding Universe.³ In particular, we are only interested in transitions where the expansion rate of the Universe is much slower than typical fluctuation time-scales in the system, *i.e.*, when $M_{Pl} \gg m$, where M_{Pl} is the Planck mass, and m is the characteristic mass scale in the model.

In this work we will study analytically the early stages of a first order phase transition. That is, we will restrict our study to temperatures above the critical temperature T_c , which is defined as the temperature at which the homogeneous part of the coarse-grained free energy density (the effective potential to some order in perturbation theory) exhibits two degenerate minima. By focusing on temperatures above the critical temperature, we will be examining the behavior of thermal fluctuations around the high temperature equilibrium state of the system. In particular, we will consider models described by a potential $V(\phi, T)$ which at high temperatures has a minimum at $\langle\phi\rangle = 0$ and that, as the temperature drops to some value T_1 , develop a new minimum at $\langle\phi\rangle = \phi_+(T)$. At the critical temperature $T_c < T_1$, $V(0, T_c) = V(\phi_+, T_c)$. We will examine thermal fluctuations

around $\langle\phi\rangle = 0$ as the temperature drops below T_1 .

The reader may be wondering why should anyone be interested in studying the kinetics of first order phase transitions above T_c . After all, it is a well-known fact that first order transitions evolve through the nucleation and subsequent percolation of bubbles larger than a critical size as the temperature drops below T_c . This is indeed the case for sufficiently strong transitions. However, for weak enough transitions the critical bubble picture must be modified, as has been observed for several condensed matter systems in the past two decades. A particularly striking example is the isotropic-nematic transition in certain liquid crystals; the transition is first order (there is a discontinuous jump of the order parameter) even though there is no release of latent heat.⁴ In fact, large amplitude fluctuations of the nematic phase within the isotropic phase have been observed *above* the critical temperature for the transition, a typical example of the so-called pre-transitional phenomena in condensed matter literature.⁵ (Interestingly enough, the free energy density used to describe the isotropic-nematic transition, the Landau-de Gennes free energy, is formally identical to the effective potential obtained for the electroweak transition.) Here, we would like to investigate this possibility within the context of hot field theories. The point is that the usual vacuum decay mechanism for first order transitions relies on results obtained within homogeneous nucleation calculations, which assume that below T_c the system has small fluctuations about the metastable phase. In other words, it is assumed that the path integral controlling the transition rate is dominated by its saddle point, given by the solution to the Euclidean equations of motion. Small fluctuations are then incorporated by evaluating the path integral by a Gaussian approximation. But as the transition grows weaker, large amplitude fluctuations about equilibrium will become more probable and the approximations used will break down.⁶ Instead of expanding about a homogeneous metastable background, one should be expanding about an inhomogeneous background consisting of these large amplitude sub-critical fluctuations. This is clearly

a very hard task which nevertheless must be undertaken if we are to better understand the dynamics of weak first order transitions. The present work is but a first step in this direction as we attempt to obtain the equilibrium distribution of sub-critical bubbles at temperatures above T_c . This way we can obtain the fraction of the total volume which is occupied by the broken symmetric phase as T_c is approached from above, and thus examine the validity of the homogeneous nucleation picture as a function of the strength of the phase transition. Gleiser, Kolb and Watkins⁷ (GKW) have first suggested this possibility, concentrating on correlation volume bubbles, but they did not take into account that subcritical bubbles are unstable and, therefore, shrink. (See, however, their comments in Section 3.3). Later, Anderson⁸ remarked that subcritical bubbles may disappear not only by shrinking but also due to thermal noise. As a result of any of these two mechanisms he concluded that the fraction of the total volume filled by bubbles of the unstable phase will be much smaller than proposed by GKW. Here we incorporate these two disappearance mechanisms into a kinetic description of the transition in order to investigate the importance of sub-critical bubbles near the critical temperature. We find that sub-critical fluctuations become progressively more important as the strength of the transition weakens, until the point where the approximations we use in order to treat the problem analytically break down and our analysis cannot be trusted quantitatively. However, we believe that our results are indicative of the relevance of pre-transitional phenomena in the study of sufficiently weak phase transitions in hot field theories.

In the next Section we will obtain the equation governing the kinetics of sub-critical bubbles for temperatures just below T_1 . In Section 3 we obtain the expressions for the thermal nucleation rates needed to solve the kinetic equation. In Section 4 we solve the kinetic equation in three regimes; assuming that the shrinking dominates the disappearance of subcritical bubbles, assuming that thermal noise dominates, and finally neglecting both shrinking and thermal noise. Starting with the whole volume occupied

only by the stable phase, we obtain the equilibration time scale and the equilibrium number density of bubbles of the unstable phase in the three regimes at the particular temperature considered, and establish the conditions for each of the three processes to dominate the kinetics. In Section 5 we compare the three time scales in the context of the standard electroweak model. As expected, for strong enough transitions shrinking or thermal noise dominate and the sub-critical bubbles play a negligible role during the phase transition. As the transition weakens we find that a larger fraction of the total volume is occupied by the broken-symmetric phase. The approximations we used to analytically solve the kinetic equation break down when a large fraction of the volume is occupied by sub-critical bubbles of the broken-symmetric phase, and we cannot carry our study into the limit of very weak transitions. But our results clearly suggest that for weak enough transitions a departure of the usual vacuum decay mechanism is to be expected. Concluding remarks are presented in Section 6.

2. KINETIC EQUATION

Let $n(R, t)$ be the number density of sub-critical bubbles of radius R at time t . Thus, $\frac{\partial n(R, t)}{\partial R} dR$ is the number per unit volume of bubbles of radii between R and $R + dR$. Since the bubbles can shrink, their radius R is a function of time, $R(t)$. We will only consider bubbles with $R \geq \xi$, where ξ is the correlation length for fluctuations around equilibrium, given by

$$\xi^{-2}(T) = V''(\phi = 0, T). \quad (2.1)$$

Bubbles with $R \sim \xi$ will be statistically dominant since any larger fluctuation has larger free-energy and is exponentially suppressed. (Recall that for $T \geq T_c$ the free energy of bubbles is a monotonically increasing function of R .⁷⁾

Consider a large volume V filled by the stable phase $\langle\phi\rangle = 0$, which is cooled down from high temperatures to a temperature just below T_1 . Bubbles with radius R of the new phase with $\langle\phi\rangle = \phi_+$ will be thermally nucleated in the background of the phase $\langle\phi\rangle = 0$ with number density $n(R, t)$; the bubbles are energetically unfavored and will shrink away with some velocity dR/dt . Shrinking will always be present, unless there is a stabilizing mechanism for the bubbles, as in non-topological solitons.⁹ [We would like to stress though that not much is known about nonlinear bubble collapse. Naively, one would expect small, unstable bubbles to collapse in a time $\sim R$. In fact, recent results on the evolution of unstable bubbles found a remarkable *enhancement* of the lifetime of bubbles by 3 to 4 orders of magnitude, as long as their initial radius is larger than about 2.5ξ and their amplitude at the center probes the nonlinearity of the potential.¹⁰] Thermal noise may also destroy small sub-critical bubbles. The importance of this effect will depend both on the ratio of bubble size to thermal length (R/T^{-1}), and on the strength of the coupling of the bubbles with the thermal background, which we parameterize with a dimensionless coefficient a . Anderson⁸ wrote this “thermal destruction” rate as $\simeq aT(RT)^3$. This expression would imply that the rate of disappearance of a bubble due to thermal noise increases with its size, a result we find counter-intuitive. Instead, we will conservatively take this rate to be of order aT , and thus independent of bubble size. As we are mostly interested in small bubbles here, we expect that this overestimate of the true rate will not compromise our results. In any case, it will only make our final results stronger, as the final fraction of volume occupied by sub-critical bubbles will be larger than what we obtain. This expression yields a lower rate than that of Ref. 8, since $T > R^{-1}$. We take, therefore, the rate per unit volume of disappearance of sub-critical bubbles due to thermal noise to be $\Gamma_{TN}(R) \simeq aT/\frac{4}{3}\pi R^3$.

We now proceed to obtain the rate equation. We can say, quite generally, that the number of bubbles that at time $t + dt$ have radius $R + dR$ in a volume $V \gg \xi$ is equal

to the number of bubbles at time t with radius R plus the net change in the number of bubbles in the range $(R, R + dR)$ due to thermal nucleation, disappearance due to thermal noise and shrinking in the time interval dt ,

$$\begin{aligned} n(R + dR, t + dt)V = & n(R, t)V + \left(\frac{V_0}{V}\right) \Gamma_{0 \rightarrow +}(R)V dt \\ & - \left(\frac{V_+}{V}\right) \Gamma_{+ \rightarrow 0}(R)V dt - \left(\frac{V_+}{V}\right) \Gamma_{TN}(R)V dt. \end{aligned} \quad (2.2)$$

Thus, to first order in dR and dt we obtain, after subtracting $n(R + dR, t)$ to both terms and dividing by $V dt$,

$$\begin{aligned} \frac{\partial n(R, t)}{\partial t} = & -\frac{\partial n(R, t)}{\partial R} \left(\frac{dR}{dt}\right) + \left(\frac{V_0}{V}\right) \Gamma_{0 \rightarrow +}(R) \\ & - \left(\frac{V_+}{V}\right) \Gamma_{+ \rightarrow 0}(R) - \left(\frac{V_+}{V}\right) \Gamma_{TN}(R) \end{aligned} \quad (2.3)$$

Here, $\Gamma_{0 \rightarrow +}(R)$ ($\Gamma_{+ \rightarrow 0}(R)$) is the rate per unit volume for the thermal nucleation of a bubble of radius R of phase $\phi = \phi_+$ within the phase $\phi = 0$ (phase $\phi = 0$ within the phase ϕ_+). These were the only two rates considered in GKW.⁷ The volume ratios $V_{0(+)}/V$ take into account the fact that the total volume in each phase changes in time due to the evolution of $n(R, t)$. The initial conditions we choose are

$$V_0(t = 0) = V \quad \text{and} \quad V_+(t = 0) = 0 \quad , \quad (2.4)$$

that is, all volume V is initially in the phase $\phi = 0$. Also, V_+ must be understood as the volume of the $(+)$ -phase in bubbles of radius R *only*, since we are following the evolution of $n(R, t)$. Thus $\left(\frac{V_+}{V}\right) = \frac{4}{3}\pi R^3 n(R, t)$. In Eq. (2.3) bubbles of radius R can disappear due to their shrinking (accounted for by the first term in the right-hand side with $dR/dt < 0$), due to thermal noise (last term in the right-hand side), and due to the nucleation of bubbles of the (0) -phase in their interior. This latter process will, in general, involve bubbles of (0) -phase of different radii as we will discuss later on.

Because

$$\left(\frac{V_0}{V}\right) = 1 - \int_{\xi}^{\infty} dR \left[-\frac{4}{3}\pi R^3 \frac{\partial n(R, t)}{\partial R} \right] \quad , \quad (2.5)$$

Eq. (2.3) is an integro-differential equation and must be solved within certain approximations, as with any Boltzmann-like equation. In a complete treatment of the kinetics we should also write an equation for the fluctuating regions of the (0)-phase within the (+)-phase. Furthermore, there would be two contributions to V_0 , a connected background volume and a bubble-like, disconnected volume. The topology of the two-phase system will, in general, be very complicated. Thus, we choose the range of temperatures in which it is consistent to restrict the (0)-phase to the background, as will be clear later.

Before we move on, we comment on two other possible contributions to $n(R, t)$ which we will not discuss here; a) Induced nucleation: small bubbles may act as seeds for the nucleation of other bubbles in their neighborhood due to the gain in surface energy. There should be an enhancement of the nucleation rate due to the presence of small bubbles in the background, very much like the presence of impurities in condensed matter systems. b) Collision of bubbles: small bubbles may acquire a thermal velocity and collide forming larger bubbles. Of course, for very short lived bubbles none of these processes should be very important. However, we again stress that large enough bubbles can be quite long-lived, and that a truly realistic scenario of weak first order transitions will be much more complicated than our simple model. The worth of the present effort relies on it being the first attempt to go beyond GKW, by incorporating more complicated out-of-equilibrium processes in the usual description of phase transitions in field theories.

3. THERMAL NUCLEATION RATES

In order to solve Eq. (2.3) we need to determine the thermal nucleation rates $\Gamma_{0(+)\rightarrow+(0)}$. For temperatures $T \lesssim T_1$, it is reasonable to assume that most bubbles will be nucleated

with radius $R \gtrsim \xi$, where ξ was defined in Eq. (2.1). For simplicity we will assume the same correlation length for fluctuations around both minima, even though at T_1 the correlations about ϕ_+ diverge. Following GKW we write for the rates,

$$\Gamma_{0(+)\rightarrow+(0)}(R) = AT^4 \exp \left[-F(\bar{\phi}_{+(0)})/T \right] \quad , \quad (3.1)$$

where A is a constant of order unity and the ansatz for the sub-critical bubble configurations $\bar{\phi}$ is

$$\bar{\phi}_+(r) = \phi_+ \exp \left[-r^2/R^2 \right] \quad ; \quad \bar{\phi}_0(r) = \phi_+ \left(1 - \exp \left[-r^2/R^2 \right] \right) , \quad (3.2)$$

with the free energy functional $F(\bar{\phi})$ of the bubble configuration given by

$$F(\bar{\phi}) = 4\pi \int r^2 dr \left[\frac{1}{2} \left(\frac{d\bar{\phi}}{dr} \right)^2 + V(\bar{\phi}, T) \right] \quad . \quad (3.3)$$

In order to move on, we will choose a specific (but quite general) potential,

$$V(\phi, T) = \frac{m^2(T)}{2} \phi^2 - \gamma(T) \phi^3 + \frac{\lambda(T)}{4} \phi^4 \quad , \quad (3.4)$$

where $\gamma(T)$ and $\lambda(T)$ are positive definite functions of T and $m^2(T)$ can be negative below a certain temperature $T_2 < T_c$. This potential has a minimum at $\phi = 0$ as long as $m^2(T) > 0$, and a local minimum at $\phi_+ = \left[3\gamma(T) + \sqrt{9\gamma^2(T) - 4m^2(T)\lambda(T)} \right] / 2\lambda(T)$, which appears at a temperature T_1 given by the solution of $\gamma^2(T_1) = 4m^2(T_1)\lambda(T_1)/9$, when $\phi_+(T_1) = 3\gamma(T_1)/2\lambda(T_1)$. Below T_c this minimum becomes the global minimum and the minimum at $\phi = 0$ becomes metastable. Using Eqs. (3.2) and (3.4) in Eq. (3.3), we obtain for the free energy $F(\bar{\phi})$,¹¹

$$F(\bar{\phi}_{0(+)}) = \alpha_{0(+)} R + \beta_{0(+)} R^3 \quad , \quad (3.5)$$

where

$$\alpha_0 = \alpha_+ = \frac{3\sqrt{2}}{8} \pi^{3/2} \phi_+^2 \quad , \quad (3.6)$$

$$\begin{aligned} \beta_0 = & \pi^{3/2} \phi_+^2 \left[\frac{m^2(T)}{8} (\sqrt{2} - 8) + \gamma(T) \phi_+ \left(3 - \frac{3\sqrt{2}}{4} + \frac{\sqrt{3}}{9} \right) \right. \\ & \left. + \lambda(T) \phi_+^2 \left(-1 + \frac{3\sqrt{2}}{8} - \frac{\sqrt{3}}{9} + \frac{1}{32} \right) \right] , \end{aligned} \quad (3.7)$$

and

$$\beta_+ = \pi^{3/2} \phi_+^2 \left[\frac{m^2(T) \sqrt{2}}{8} - \frac{\sqrt{3}}{9} \gamma(T) \phi_+ + \frac{\lambda(T)}{32} \phi_+^2 \right] . \quad (3.8)$$

4. SOLVING THE KINETIC EQUATION

We will solve Eq. (2.3) in different regimes assuming that the dominant process for the disappearance of bubbles is: 1.) their shrinking, 2.) thermal noise and 3.) the nucleation of bubbles of the true vacuum inside them. This will allow us to make a direct comparison of the results for the relaxation time scale and equilibrium number density obtained in these regimes, in the context of the electroweak transition in the next Section.

4.1 Kinetic Equation with Shrinking

We include only the shrinking term, neglecting the last two terms of Eq. (2.3), and solve this equation with the approximation,

$$\left(\frac{V_0}{V} \right) = 1 - \int_{\xi}^{\infty} dR \left[-\frac{4}{3} \pi R^3 \frac{\partial n(R, t)}{\partial R} \right] \equiv 1 - |I| \simeq 1 . \quad (4.1)$$

In words, we will solve the kinetic equation in the regime in which the volume in the (+)-phase, given by the sum over all bubbles with radii from ξ to ∞ (actually, to $R \sim V^{1/3} \gg \xi$), is small, i.e. $|I| \ll 1$. We will check our results for consistency after obtaining $n(R, t)$. With this approximation the kinetic equation becomes,

$$\frac{\partial n(R, t)}{\partial t} = \frac{\partial n(R, t)}{\partial R} [f(R)] + \Gamma_{0 \rightarrow +}(R) , \quad (4.2)$$

where we wrote $dR/dt = -f(R) < 0$. The positive definite function $f(R)$ encompasses the physically interesting cases $dR/dt = v = \text{constant}$, and $dR/dt \sim 1/R$. We enforced the shrinking explicitly by choosing the minus sign in dR/dt . The rate $\Gamma_{0 \rightarrow +}$ was defined in Eq. (3.1), with the free energy given by Eq. (3.5) with α_+ and β_+ given by Eqs. (3.6) and (3.8) respectively. In order to solve this equation analytically we make one further approximation; we neglect the volume term in the expression for the free energy, so that the nucleation rate can be written as

$$\Gamma_{0 \rightarrow +}(R) \simeq AT^4 \exp[-\alpha_+ R/T] \equiv g(R) \quad . \quad (4.3)$$

Here we have defined the function $g(R)$. Given that most bubbles have radii $R \gtrsim \xi$ this should be a good approximation that must be tested in each application. For the electroweak case, independently of the parameters of the model, we obtain $\alpha_+ \xi / \beta_+ \xi^3 \simeq 6.65$ for $T = T_1$. The ratio increases for lower temperatures.

In order to solve the above equation, first note that the time independent equilibrium number density at the temperature chosen, $\bar{n}(R)$, is the solution of

$$\frac{\partial \bar{n}(R)}{\partial R} = -\frac{g(R)}{f(R)} \quad , \quad (4.4)$$

which is easily obtained as

$$\bar{n}(R) = \bar{n}(\xi) - \int_{\xi}^R \frac{g(R)}{f(R)} dR \quad . \quad (4.5)$$

Choosing $f(R) = v$ and the definition of $g(R)$ above we obtain,

$$\bar{n}(R) = \frac{AT^5}{v\alpha_+} \exp[-\alpha_+ R/T] \quad . \quad (4.6)$$

Using Eq. (4.4), we can rewrite the kinetic equation as

$$\frac{\partial Y(R, t)}{\partial t} = \frac{\partial Y(R, t)}{\partial R} f(R), \quad (4.7)$$

where we introduced the departure from equilibrium, $Y(R, t) \equiv n(R, t) - \bar{n}(R)$. This equation is solved by writing

$$Y(R, t) = X(R) \exp[-t/\tau_1] \quad , \quad (4.8)$$

so that τ_1 is the relaxation time and the solution approaches the equilibrium distribution as $t \rightarrow \infty$. Choosing again $f(R) = v$ we obtain upon substitution in Eq. (4.7),

$$X(R) = X_0 \exp[-R/v\tau_1] \quad , \quad (4.9)$$

where X_0 is a constant, which, together with τ_1 , will be fixed by the initial condition $Y(R, 0) = -\bar{n}(R)$. Using Eq. (4.6) we find

$$X_0 = -\frac{AT^5}{v\alpha_+} \quad \text{and} \quad \tau_1 = \frac{T}{v\alpha_+} \quad . \quad (4.10)$$

Putting the results together we find for the solution of the kinetic equation,

$$n(R, t) = \bar{n}(R) \left(1 - e^{-t/\tau_1}\right) = \frac{AT^5}{v\alpha_+} e^{-\alpha_+ R/T} \left[1 - e^{-\alpha_+ vt/T}\right] \quad . \quad (4.11)$$

Thus, the relaxation time to approach the equilibrium distribution $\bar{n}(R)$ is the time for a bubble of characteristic radius $v\tau_1 = T/\alpha_+$ to shrink with constant velocity v . (The reason why τ_1 is independent of R is due to our choice of constant shrinking speed for all bubbles.) Substituting this solution into the consistency condition given in Eq. (4.1) we obtain

$$|I| = \frac{4}{3} \pi \xi^3 \bar{n}(\xi) \left[1 + 3 \left(\frac{T}{\alpha_+ \xi}\right) + 6 \left(\frac{T}{\alpha_+ \xi}\right)^2 + 6 \left(\frac{T}{\alpha_+ \xi}\right)^3\right] \ll 1 \quad . \quad (4.12)$$

The consistency condition is then an expression of the validity of the semi-classical approximation, which requires that $\exp[-F(\bar{\phi}_+)/T] \simeq \exp[-\alpha_+ R/T] \ll 1$; if the smallest bubble has a large nucleation barrier ($= F_+/T$) its production is exponentially suppressed and only a negligible fraction of the total volume will be occupied by the (+)-phase.

4.2 Kinetic Equation with Thermal Destruction

In Ref. 8 Anderson writes the thermal destruction rate per unit volume as

$$\Gamma(R)_{TN} \simeq T^4 . \quad (4.13)$$

With this expression, the rate of disappearance of a bubble increases with its volume. As mentioned above, we think this is counter-intuitive, since larger bubbles should be less prone to be destroyed by thermal noise than smaller ones, not the contrary. In lack of a better understood expression we will take this rate to be aT , on dimensional grounds, independently of the size of the bubble. The rate per unit volume of the disappearance of bubbles due to thermal noise is then,

$$\Gamma(R)_{TN} = \frac{aT}{\frac{4\pi}{3}R^3} . \quad (4.14)$$

The constant a is proportional to the coupling of the bubble to the thermal bath to some power. This is a difficult quantity to obtain microscopically, and will depend on how the field couples to itself and to other fields in the model. There has been some progress recently in the computation of the viscosity coefficient in hot field theories, although the results are model dependent.¹² However, it is by now clear that viscosity (which is related to the coupling to the bath by the fluctuation-dissipation theorem) in field theories is at least a two-loop effect.

The kinetic equation is now,

$$\frac{\partial n(R, t)}{\partial t} = \left(\frac{V_0}{V} \right) \Gamma_{0 \rightarrow +}(R) - \frac{4}{3} \pi R^3 n(R, t) \Gamma_{TN}(R) . \quad (4.15)$$

We will again make the approximation of Eq. (4.1). Using Eqs. (4.3) and (4.13), the kinetic equation becomes

$$\frac{\partial n(R, t)}{\partial t} = g(R) - aTn(R, t) . \quad (4.16)$$

Thus, the time independent equilibrium number density is now

$$\bar{n}(R) = \frac{g(R)}{aT} = \frac{AT^3}{a} e^{-\alpha_+ R/T} , \quad (4.17)$$

and we solve the equation for $(n - \bar{n})$ with the ansatz

$$n(R, t) - \bar{n}(R) = X(R)e^{-t/\tau_2} \quad , \quad (4.18)$$

and the initial condition $n(R, 0) = 0$, which yields $X(R) = \bar{n}(R)$. Thus, $\tau_2^{-1} = aT$ and

$$n(R, t) = \bar{n}(R)(1 - e^{-aTt}) = \frac{AT^3}{a}e^{-\alpha_+R/T}(1 - e^{-aTt}) \quad (4.19)$$

The relaxation time is thus $\tau_2 = (aT)^{-1}$ as expected. Comparing this value with the relaxation time when shrinking alone is considered $\tau_1 \simeq T/\alpha_+v$, as obtained in the previous section, we see that, under the approximations we choosed, thermal noise will be dominant, *i.e.* $\tau_2 < \tau_1$, for

$$a > \frac{\alpha_+v}{T^2} \simeq \frac{\phi_+^2 v}{T^2}. \quad (4.20)$$

Here we have taken $\alpha_+ \simeq \phi_+^2$ from Eq. (3.6). This is an important condition, since, as we mentioned above, a is proportional to some power of the coupling of the field ϕ with the thermal bath, and ϕ_+/T is a measure of the strength of the transition. (For successful baryogenesis during the electroweak scale one needs roughly $\phi_+/T \gtrsim 1$.¹³) Thus, thermal noise only dominates the dynamics over shrinking for sufficiently weak transitions, *and* for strong enough coupling to the thermal bath. The consistency condition in Eq. (4.1) is again given by Eq. (4.12), but in this case $\bar{n}(\xi)$ is given by Eq. (4.17). Thus, this condition is again an expression of the validity of the semiclassical approximation, as explained below Eq. (4.17).

4.3 Kinetic Equation Without Shrinking and Thermal Destruction

Neglecting the shrinking of bubbles and their coupling to thermal noise, the kinetic equation is

$$\frac{\partial n(R, t)}{\partial t} = \left(\frac{V_0}{V}\right) \Gamma_{0 \rightarrow +}(R) - \left(\frac{V_+}{V}\right) \Gamma_{+ \rightarrow 0} \quad , \quad (4.21)$$

where the nucleation rates have been defined before. In order to be consistent with our previous approach, we will again take $V_0/V \simeq 1$. Since $(V_+/V)\Gamma_{+\rightarrow 0}(R)$ is the rate at which bubbles of $(+)$ -phase of radius R disappear due to the nucleation of a region of (0) -phase in its interior, as mentioned before, V_+ is the total volume of all bubbles of radius R (and not the total volume of $(+)$ -phase). Thus we have $V_+/V = (4\pi/3)R^3 n(R, t)$. Consistency with the assumption $V_0/V \simeq 1$ as $t \rightarrow \infty$ requires that $(4\pi/3)R^3 \bar{n}(R) \ll 1$. Since most bubbles of the $(+)$ -phase will have $R \simeq \xi$, this condition is essentially the same as that of Eq. (4.12), as it should.

In principle, to obtain $\Gamma_{+\rightarrow 0}$ we should sum over bubbles of the (0) -phase of different radii which can trigger the disappearance of regions of the $(+)$ -phase. This sum should be dominated by the contribution of bubbles of radius ξ . In this case the dominant process for wiping out regions of $(+)$ -phase is nucleation of correlation volume regions of the (0) -phase, as was explicitly assumed in the work of GKW. However, whenever we will need to write $\Gamma_{+\rightarrow 0}$ explicitly to obtain analytic expressions, we will approximate it with the rate of thermal nucleation of bubbles of radius R , $\Gamma_{+\rightarrow 0} \simeq \Gamma_{+\rightarrow 0}(R)$. We expect this approximation to be good since most bubbles of $(+)$ -phase have radii R not much larger than ξ anyway. The kinetic equation is, therefore,

$$\frac{\partial n(R, t)}{\partial t} = \Gamma_{0 \rightarrow +}(R) - \frac{4\pi}{3} R^3 n(R, t) \Gamma_{+\rightarrow 0} \quad . \quad (4.22)$$

We can rewrite it as,

$$\frac{\partial z(R, t)}{\partial t} = -q(R)z(R, t), \quad z(R, t) \equiv p(R) - q(R)n(R, t) \quad , \quad (4.23)$$

where we defined

$$p(R) \equiv \Gamma_{0 \rightarrow +}(R) \quad \text{and} \quad q(R) \equiv \frac{4\pi}{3} R^3 \Gamma_{+\rightarrow 0} \quad . \quad (4.24)$$

The solution to Eq. (4.23) is simply $z(R, t) = z_0 \exp[-q(R)t]$, or, using the initial condition

$$n(R, 0) = 0,$$

$$n(R, t) = \bar{n}(R) [1 - e^{-q(R)t}] \quad ; \quad \bar{n}(R) = \frac{p(R)}{q(R)} . \quad (4.25)$$

For a given radius R , the relaxation time is then

$$\tau_3(R) = [q(R)]^{-1} = \left[\frac{4\pi}{3} R^3 \Gamma_{+\rightarrow 0} \right]^{-1} , \quad (4.26)$$

that is, the time scale for the nucleation of a bubble of the (0)–phase inside a bubble of radius R of the (+)–phase. Note that τ_3 depends exponentially on R^3 ; bubbles of larger radii will approach the equilibrium distribution at a much slower rate than smaller bubbles. The fastest bubbles to equilibrate are those with $R \simeq \xi$. With the approximation $\Gamma_{+\rightarrow 0} \simeq \Gamma_{+\rightarrow 0}(R)$ and using the expressions of Section 2 for the nucleation rates, the equilibrium distribution is given by

$$\frac{4\pi}{3} R^3 \bar{n}(R) = \exp \left[-\frac{(\beta_+ - \beta_0) R^3}{T} \right] . \quad (4.27)$$

The consistency condition then states that the volume occupied by the (+)–phase is small,

$$\frac{4\pi}{3} \xi^3 \bar{n}(\xi) = \exp \left[-\frac{(\beta_+ - \beta_0) \xi^3}{T} \right] \ll 1 . \quad (4.28)$$

5. APPLICATION TO THE ELECTROWEAK TRANSITION

For a given model with a first order transition which of the approaches above better describes the dynamics of thermal fluctuations? We are limited to study the transition at $T \lesssim T_1$ when only small sub-critical fluctuations exist. However, even for this limited temperature range we should be able to examine the importance of incorporating the shrinking of the bubbles and their coupling to thermal noise into the description of the kinetics. In this Section we will do this in the context of the standard electroweak model

using the 1-loop approximation to the effective potential. Even though recent work has shown that the 1-loop approximation is not adequate due to large infrared corrections at T_c ,¹⁴ we will take the 1-loop potential as an example of how to apply our methods. Application to other potentials is quite straightforward.

We will compare the different relaxation time scales obtained above, τ_1 , τ_2 and τ_3 , and investigate which mechanism for the suppression of regions of broken-symmetric phase will dominate as the strength of the transition changes. Our strategy is as follows. First we compare τ_1 with τ_2 , that is, we compare the relaxation time scale incorporating only shrinking (Section 4.1) with the relaxation time scale incorporating only thermal noise (Section 4.2). We have already shown under our assumptions that thermal noise dominates over shrinking if the parameter a satisfies the inequality $a > \alpha_+ v/T^2$, Eq. (4.20). Since a is a free parameter in our model we have two possibilities, depending on the value of a . If a does not satisfy the inequality, shrinking dominates and thermal noise is a sub-dominant mechanism during the approach to equilibrium. Otherwise, shrinking is the sub-dominant mechanism. Since we then will know the values of a for which shrinking or thermal noise dominates as a function of the strength of the transition, we can proceed by comparing these two processes with the inverse thermal nucleation rate time scale, given by τ_3 (Section 4.3). As a result, we will be able to establish which mechanism dominates the dynamics as the transition's strength varies for different values of a .

We take the Higgs mass to be the parameter that controls the strength of the transition, while we fix the top mass at $m_T = 130$ GeV. For the 1-loop approximation to the electroweak potential, we can write¹⁵

$$V_{\text{EW}}(\phi, T) = D (T^2 - T_2^2) \phi^2 - ET\phi^3 + \frac{\lambda_T}{4} \phi^4 \quad , \quad (5.1)$$

where the constants D and E are given by $D = [6(m_W/\sigma)^2 + 3(m_Z/\sigma)^2 + 6(m_T/\sigma)^2] / 24$,

and $E = [6(m_W/\sigma)^3 + 3(m_Z/\sigma)^3]/12\pi \simeq 10^{-2}$. T_2 is the temperature at which the origin becomes unstable, given by

$$T_2 = \sqrt{(m_H^2 - 8B\sigma^2)/4D} \quad , \quad (5.2)$$

where the physical Higgs mass is given in terms of the 1-loop corrected λ as $m_H^2 = (2\lambda + 12B)\sigma^2$, with $B = (6m_W^4 + 3m_Z^4 - 12m_T^4)/64\pi^2\sigma^4$. We use $m_W = 80.6$ GeV, $m_Z = 91.2$ GeV, and $\sigma = 246$ GeV. The temperature corrected Higgs self-coupling is

$$\lambda_T = \lambda - \frac{1}{16\pi^2} \left[\sum_B g_B \left(\frac{m_B}{\sigma} \right)^4 \ln(m_B^2/c_B T^2) - \sum_F g_F \left(\frac{m_F}{\sigma} \right)^4 \ln(m_F^2/c_F T^2) \right], \quad (5.3)$$

where the sum is performed over bosons and fermions (in our case only the top quark) with their respective degrees of freedom $g_{B(F)}$. Also, $\ln c_B = 5.41$ and $\ln c_F = 2.64$.

This potential is equivalent to the potential of Eq. (3.4), with the replacements $m^2(T) = 2D(T^2 - T_2^2)$, $\gamma(T) = ET$, and $\lambda(T) = \lambda_T$. The temperatures T_1 and T_c are given by

$$T_1 = T_2/\sqrt{1 - 9E^2/8\lambda_T D}, \quad \text{and} \quad T_c = T_2/\sqrt{1 - E^2/\lambda_T D} \quad . \quad (5.4)$$

From these values for T_1 and T_c , it is clear that the quantity $x \equiv E^2/\lambda_T D$ can be used as a measure of the strength of the transition. For $x = 0$, the transition is second order. As shown in Ref. 17, $x \simeq 0.027$ at $m_H = 60$ GeV, becoming smaller for larger Higgs masses. The transition is already quite weak at the experimental lower bound of $m_H = 57$ GeV.¹⁶ In Fig. 1 we show the quantity x as a function of the Higgs mass for $m_H \geq 35$ GeV.

With the potential above, we can easily compute the nucleation rates using the expressions of Section 2, as done by Gleiser and Kolb.¹¹ We have then all that is needed to compute the relaxation time scales τ_1 , τ_2 , and $\tau_3(\xi)$ [since τ_3 depends on R , we consider only the fastest bubbles to equilibrate, those of radius $\sim \xi$], and the equilibrium distributions $\bar{n}(R)$.

Let us start by comparing the relaxation time scales for the shrinking term and the thermal destruction term, following Eq. (4.20). In Fig. 2 we plot $\alpha_+(T_1)/T_1^2$ (we took $v = 1$) as a function of the Higgs mass. Thermal noise dominates over shrinking if a lies above the curve, for a given Higgs mass. Notice that for small Higgs masses, *i.e.* strong transitions, shrinking should be the dominant process unless the coupling to the bath is unrealistically large (or non-perturbative). However, since there are uncertainties in the precise expression for the thermal destruction rate, we will consider both possibilities, with shrinking and thermal noise dominating the dynamics, respectively. If shrinking dominates, we can neglect thermal noise and compare τ_1 and $\tau_3(\xi)$ directly. This is done in the continuous lines of Fig. 3 where we show the ratio of $\tau_1/\tau_3(\xi)$ as a function of the Higgs mass. The results are shown for two different shrinking velocities, $v = 1$ and $v = 0.4$. For both velocities, with Higgs masses below 52 GeV we find that $\tau_1 \ll \tau_3(\xi)$, so that the approach to equilibrium is dominated by the shrinking. In Fig. 4 we show the two consistency conditions Eqs. (4.12) and (4.28) as a function of the Higgs mass. We see that the fraction of the total volume occupied by the (+)-phase increases rapidly as the Higgs mass increases. That is, as the transition grows weaker a larger fraction of the volume is occupied by the broken-symmetric phase. Even though our results indicate that we cannot trust our approximations for $m_H \gtrsim 53$ GeV, where $V_+/V \gtrsim 10\%$, it is quite clear that the reason our approximations break down is precisely the large fraction of the volume occupied by sub-critical bubbles. These results suggest that pre-transitional phenomena will be present for weak enough transitions and that we should expect modifications of the vacuum decay picture in this case. A precise description of the transition is beyond the scope of the present formalism.

We now consider the case in which thermal noise dominates over shrinking. By inspecting Fig. 2, we can find an appropriate value of a which is large enough for the inequality Eq. (4.20) to be always satisfied. We will take $a = 0.5$ as an illustration, even

though this value is probably unrealistically large. In Fig. 3 the dashed line shows the ratio $\tau_2/\tau_3(\xi)$ as a function of the Higgs mass. Note that again thermal noise becomes sub-dominant as the Higgs mass is increased, although it does so at a slower rate than the shrinking term. However, the qualitative conclusion is the same in both cases. For large enough Higgs masses, the fraction of the total volume in the broken-symmetric phase becomes substantial due to the increasing weakness of the transition. This weakness is characterized by the fact that both the shrinking of sub-critical bubbles as well as their destruction due to thermal noise become less important in the description of the transition. Since the breakdown of our approximations is due to the failure of the semi-classical, or dilute gas approach, we are confident that as the Higgs mass increases beyond the limit of validity of our approximations a regime will be reached in which a departure from the usual false vacuum decay mechanism is to be expected, although at this point we cannot afford to make a quantitative prediction. A rough estimate may be obtained by comparing, at the nucleation temperature, the typical distance between sub-critical bubbles to the radius of a critical bubble obtained by the usual calculations. If the distance between sub-critical bubbles is of the order of the critical radius, the usual nucleation mechanism must be revised. Work on this topic is currently in progress.

6. CONCLUDING REMARKS

We have obtained a kinetic equation describing the approach to equilibrium in first-order transitions for temperatures above the critical temperature. For sufficiently strong transitions and making approximations that we consider reasonable, we were able to analytically solve the kinetic equation for three different regimes, determined by the dominant mechanism responsible for the suppression of regions of the broken-symmetric phase within the symmetric phase. The three processes are the shrinking of the sub-

critical bubbles, their destruction due to the thermal bath to which they couple, and by thermal nucleation of regions of the symmetric phase in their interior. By obtaining the relaxation time-scales in all regimes, we were able to study the relative importance of each process in the early stages of the transition.

We applied our approach to the standard electroweak transition, showing that for Higgs masses below 55 GeV or so, the approach to equilibrium is dominated by either shrinking or thermal destruction, depending on the strength of the coupling of bubbles to the thermal bath. In this case, the total volume occupied by the equilibrium distribution of sub-critical bubbles of the broken-symmetric phase is negligible, and the transition proceeds by the usual vacuum decay mechanism. As the Higgs mass increases and the transition becomes progressively weaker, we found that a larger fraction of the total volume becomes occupied by the broken-symmetric phase, forcing the breakdown of our analytical approximations. However, it is clear from our results that a regime will eventually be reached in which a substantial fraction of the volume is in the broken-symmetric phase as the critical temperature is reached. In this case we should expect a departure from the usual vacuum decay mechanism. We cannot be quantitative about the value of the Higgs mass (for a given top mass) where this occurs due to the limitations of our analytical approach. Given that the lower bound on the Higgs mass is now above 60 GeV, we expect very interesting physics to be lurking behind our present knowledge of the dynamics of weak first-order transitions.

ACKNOWLEDGEMENTS

We thank E. W. Kolb and D. Seckel for useful discussions. We would like to thank the Institute for Theoretical Physics at Santa Barbara for its hospitality during the Cosmo-

logical Phase Transitions program, where this work was initiated. (GG) was supported in part by the Department of Energy under the contract No. DE-FG03-91ER 40662, Task C. (MG) was supported in part by National Science Foundation grants No. PHY-9204726 at Dartmouth and No. PHY-8904035 at the Institute for Theoretical Physics at Santa Barbara.

References

1. For reviews see, J. D. Gunton, M. San Miguel, and P. S. Sahni, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, London, 1983), Vol. 8; J. S. Langer, Lectures presented at l'École de Physique de la Matière Condensée, Beg Rohu, France, August 1989, ITP preprint NSF-ITP-90-221i.
2. See the review by J. Langer in Ref. 1.
3. E.W. Kolb and M.S. Turner, *The Early Universe*, (Addison-Wesley, 1990); A.D. Linde, *Particle Physics and Inflationary Cosmology*, (Harwood Academic Publishers, 1990).
4. T. Stinton III and J. Lister, *Phys. Rev. Lett.* **25**, 503 (1970). H. Zink and W.H. de Jeu, *Mol. Cryst. Liq. Cryst.* **124**, 287 (1985). For other condensed matter systems exhibiting pre-transitional phenomena see, for example, L.A. Fernández, J.J. Ruiz-Lorenzo, M. Lombardo, and A. Tarancón, *Phys. Lett. B* **277**, 485 (1992); A. Gonzalez-Arroyo, M. Okawa, and Y. Shimizu, *Phys. Rev. Lett.* **60**, 487 (1988).

5. J.J. Stankus, R. Torre, C.D. Marshall, S.R. Greenfield, A. Sengupta, A. Tokmakoff, and M.D. Fayer, *Chem. Phys. Lett.* **194**, 213 (1992).
6. M. Gleiser and R.O. Ramos, *Phys. Lett. B* **300**, 271 (1993).
7. M. Gleiser, E. W. Kolb, and R. Watkins, *Nucl. Phys.* **B364**, 411 (1991); M. Gleiser, *Phys. Rev.* **D42**, 3350 (1990).
8. G. Anderson, *Phys. Lett. B* **295**, 32 (1992); See also, M. Dine, R. Leigh, P. Huet, A. Linde, and D. Linde, *Phys. Rev. D* **46**, 550 (1992).
9. J. Frieman, G. Gelmini, M. Gleiser, and E. W. Kolb, *Phys. Rev. Lett.* **60**, 2101 (1988).
10. M. Gleiser, Dartmouth College report No. DART-HEP-93/05, August 1993; V.G. Makhankov, *Phys. Rep. C* **35**, 1 (1978).
11. M. Gleiser and E. W. Kolb, *Phys. Rev. Lett.* **69**, 1304 (1992).
12. A. Hosoya and M. Sakagami, *Phys. Rev. D* **29**, 2228 (1984); A. Hosoya, M. Sakagami, and M. Takao, *Ann. Phys. (NY)* **154**, 229 (1984); M. Morikawa, *Phys. Rev. D* **33**, 3607 (1986).
13. A.G. Cohen, D.B. Kaplan, and A.E. Nelson, UC San Diego report No. UCSD-PTH-93-02, to appear in *Ann. Rev. Nucl. Part. Sci.*, **43**.
14. G. Boyd, D. Brahm and S.D.H. Hsu, Harvard preprint No. HUTP-92-A027 (1992), in press *Phys. Rev. D*; M. E. Carrington, *Phys. Rev. D* **45**, 2933 (1992); M. Dine *et. al.* in Ref. 8; P. Arnold, *Phys. Rev.* **D46**, 2628 (1992); M. Gleiser and E. W. Kolb, *Phys. Rev. D* **48**, 1560 (1993).
15. G.W. Anderson and L.J. Hall, *Phys. Rev.* **D45**, 2685 (1992).

16. ALEPH, DELPHI, L3 and OPAL Collaborations, as presented by M. Davier, Proceedings of the International Lepton-Photon Symposium and Europhysics Conference on High Energy Physics, eds. S. Hegerty, K. Potter and E. Quercigh, November 1991, to appear.
17. M. Gleiser and E. W. Kolb, in Ref. 14.

FIG. 1. The parameter $x = E^2/\lambda_T D$ as a function of the Higgs mass for several values of the top mass.

FIG. 2 The ration of relaxation time-scales for shrinking and thermal destruction processes as a function of the Higgs mass. The region above the curve defines the domain for thermal destruction domination.

FIG. 3. The continuous lines denote the ratio between the shrinking and reverse nucleation time-scales, as a function of the Higgs mass for shrinking velocities $v = 1$ and $v = 0.4$. The dashed line denotes the ratio between thermal destruction and reverse nucleation time-scales with parameter $a = 0.5$.

FIG. 4. The consistency conditions of Eqs. 4.12 and 4.28 as a function of the Higgs mass.